

Sharon M. Tomaski, M.D.

7720 South Broadway, Suite 480
Littleton, CO 80122
tel. 303.347.0800 fax. 303.347.1140
www.drTomaski.com

Ear Infection (Otitis Media)

What is an ear infection?

An ear infection is a bacterial infection of the middle ear (the space behind the eardrum). It usually is a complication of a cold, occurring after the cold blocks off the eustachian tube (the passage connecting the middle ear to the back of the throat). Your child's ear is painful because trapped, infected fluid puts pressure on the eardrum, causing it to bulge. Other symptoms are irritability and poor sleep.

Most children will have at least one ear infection, and over one fourth of these children will have repeated ear infections. In 5% to 10% of children, the pressure in the middle ear causes the eardrum to rupture and drain a yellow or cloudy fluid. This small hole usually heals over the next week. Children are most likely to have ear infections between the ages of 6 months and 2 years, but they continue to be a common childhood illness until the age of 8 years.

If the following treatment is carried out your child should be fine. Permanent damage to the ear or to the hearing is very rare.

How can I take care of my child?

- **Antibiotics** Take the antibiotics as your doctor has instructed. Try not to forget any of the doses. If your child goes to school or a baby sitter, arrange for someone to give the afternoon dose. If the medicine is a liquid, store the antibiotic in the refrigerator and use a measuring spoon to be sure that you give the right amount. Give the medicine until the bottle is empty or all the pills are gone. (Do not save the antibiotic for the next illness because it loses its strength.) Even though your child will feel better in a few days, give the antibiotic until it is completely gone. Finishing the medicine will keep the ear infection from flaring up again.
- **Pain relief** Acetaminophen or ibuprofen can be used to help with the earache or fever over 102°F (39°C) for a few days until the antibiotic takes effect. These medications usually control the pain within 1 to 2 hours. Earaches tend to hurt more at bedtime. To help ease the pain, you can put an ice bag or ice wrapped in a wet washcloth over the ear. This may decrease the swelling and pressure inside. Some physicians recommend a heating pad instead. Remove the cold or heat in 20 minutes to prevent frostbite or a burn.
- **Restrictions** Your child can go outside and does not need to cover the ears. Swimming is permitted as long as there is no perforation (tear) in the eardrum or drainage from the ear. Children with ear infections can travel safely by aircraft if they are taking antibiotics. Also give them an appropriate dose of ibuprofen 1 hour before take-off for any discomfort they might have. Most will not have an increase in their ear pain while flying. While coming down in elevation during an airline flight or a trip from the mountains, have your child swallow fluids, suck on a pacifier, or chew gum. Your child can return to school or day care when he or she is feeling better and the fever is gone. Ear infections are not contagious.
- **Ear recheck** Your physician will schedule a return appointment in 2 to 3 weeks for your child. At that visit, the eardrum will be looked at to be certain that the infection is cleared up and more

treatment isn't needed. Your physician may also want to test your child's hearing. Follow-up exams are very important, particularly if the infection has caused a hole in the eardrum.

How can I help prevent ear infections?

If your child has recurrent ear infections, it's time to look at how you can prevent some of them. The following list includes ways you can help your child prevent another ear infection. If some of the following items apply to your child, try to use them or talk to your health care provider about them.

- **Protect your child from second-hand tobacco smoke.** Passive smoking increases the frequency and severity of infections. Be sure no one smokes in your home or at day care.
- **Reduce your child's exposure to colds during the first year of life.** Most ear infections start with a cold. Try to delay the use of large day care centers during the first year by using a sitter in your home or a small home-based day care. Frequent hand washing can reduce the possibility of catching and passing on cold viruses.
- **Breast-feed your baby during the first 6 to 12 months of life.** Antibodies in breast milk reduce the rate of ear infections. If you're breast-feeding, continue. If you're not, consider it with your next child.
- **Avoid bottle propping.** If you bottle-feed, hold your baby at a 45° angle. Feeding in the horizontal position can cause formula and other fluids to flow back into the eustachian tube. Allowing an infant to hold his own bottle also can cause milk to drain into the middle ear. Weaning your baby from a bottle between 9 and 12 months of age will help stop this problem.
- **Control allergies.** If your infant has continuous nasal secretions, consider an allergy as a contributing factor to the ear infections, especially if your child has other allergies such as eczema.
- **Check the adenoids.** If your toddler constantly snores or breaths through his mouth, he may have large adenoids. Large adenoids can contribute to ear infections. Talk to your physician about this.

When should I call my child's health care provider?

Call IMMEDIATELY if:

- Your child develops a stiff neck.
- Your child acts very sick.

Call during office hours if:

- The fever or pain is not gone after your child has taken the antibiotic for 48 hours.
- You have other questions or concerns.

Published by McKesson Provider Technologies.



This content is reviewed periodically and is subject to change as new health information becomes available. The information is intended to inform and educate and is not a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional.

Written by B.D. Schmitt, M.D., author of "Your Child's Health," Bantam Books.

Copyright © 2005 McKesson Corporation and/or one of its subsidiaries. All Rights Reserved.

Special Instructions:

Ear



Copyright © 2002 McKesson Health Solutions, LLC. All rights reserved.